

Public health laws for prevention and control of Tuberculosis in India

Ashutosh Pathak^{1*}, Uday Gupta¹, Malik Jameel¹, Divyaraj singh¹, Mohit Kumar², Neelam Painuly³, Abhishek Agrawal⁴

¹Department of Pharmacy Practice, Teerthanker Mahaveer College of Pharmacy, Teerthanker, Mahaveer University, Moradabad, UP 244001, India.

²Department of Pharmaceutics, Teerthanker Mahaveer College of Pharmacy, Teerthanker, Mahaveer University, Moradabad, UP 244001, India. mohitgoyal21111@gmail.com; 0009-0001-8236-7396.

³Associate Professor, School of Pharmacy & Research, Dev Bhoomi Uttarakhand University, Naugaon, Dehradun, India;

Email ID : neelu.painuly@gmail.com

⁴Associate Professor, IITM COLLEGE OF PHARMACY, Murthal, Sonapat;

Email ID : abhishek060688@gmail.com

Corresponding Author:

Ashutosh Pathak

Assistant Professor, Teerthanker Mahaveer College of Pharmacy, Teerthanker Mahaveer University, Moradabad-244001, U.P., India;

Email ID : ashutosh4088@gmail.com (0009000091589356).

ABSTRACT

This paper aims to discuss the emergence, public health, and legal-regulatory aspects of tuberculosis prevention and control in India. Even though tuberculosis (TB) is considered a preventable and curable disease, it still remains one of the country's challenging public health issues. The high incidence rate, social stigma, poverty, poor nutritional balance, and overpopulation contribute to the complexity of the disease's control in India. India's TB control strategy has been changing and developing throughout time, starting from the initial period when little was known about the disease, resulting in ineffective measures, such as BCG vaccination and establishing the National TB programme in 1962. Later, a more comprehensive approach, such as the Revised National Tuberculosis Control Programme (RNTCP) based on the DOTS strategy, was introduced. Eventually, the National Tuberculosis Elimination Programme (NTEP) was developed.

The article discusses the current state of tuberculosis and the existing efforts to mitigate it. It stipulates that the problem of tuberculosis control in the community calls for a shift from an isolated set of clinical practices to a more integrated, multi-faceted public health response. The main components of this response are as follows: Early case detection and prompt treatment of drug-resistant tuberculosis. Enhanced surveillance and mandatory reporting to public and private facilities. Implementation of person-centered care in relation to population health status, access to care, treatment engagement, and support. TB prevention with an emphasis on intensified case-finding and tuberculosis preventive therapy. The One Health approach which underlines the complex interplay between human, animal, and environmental health in relation to zoonotic diseases.

Moreover, the paper outlines the principle of public health measures which aims at protecting communities while respecting human rights, privacy, and dignity to avoid stigmatization. In order to achieve the ultimate goal of eradicating the disease in India, it is essential to complement legal and policy measures with the NTEP interventions in order to address the limitations of the latter..

Keywords: Tuberculosis; Public Health Laws; TB Control; India; National Tuberculosis Elimination Programme; Disease Notification and Surveillance; Patient Rights; Legal and Regulatory Framework.

1. INTRODUCTION


Tuberculosis (TB) continues to represent among India's most enduring public-health obstacles even though it can be prevented, diagnosed and cured. Produced by *Mycobacterium tuberculosis*, this illness mainly impacts the lungs yet might also engage additional organs[1] . Its airborne method of spreading, paired with crowding, lack of money, poor nutrition, late action seeking care, insufficient knowledge and societal shame, enables ongoing spread inside neighborhoods [2] . India holds a large share of worldwide TB load, rendering prevention and management a top national goal. The historical record regarding tuberculosis management within India reveals a slow transition away from small-scale, facility-focused actions toward an organized country-wide initiative[3] . Launching the Revised National Tuberculosis Control Programme alongside the Directly Observed Treatment, Short-course method improved how cases were found, made therapy consistent, and ensured patients followed their regimens[4] . Later work evolved into the National Tuberculosis Elimination Programme (NTEP), utilizing a broad method that included spotting illness early, providing cost-free care, handling resistant drugs, offering nutrition aid, working with private providers, and engaging local populations. However, medical interventions alone are insufficient to eliminate TB[5] . Effective prevention and control also require a strong legal and policy framework that defines governmental responsibilities, protects public health, regulates healthcare services, supports disease notification and surveillance, and safeguards the rights and dignity of people affected by TB[6] . Public health laws are therefore important instruments for addressing the social and environmental determinants of transmission, ensuring access to timely diagnosis and treatment, regulating infection-prevention practices and coordinating action across health, nutrition, housing, labour and local-government sectors[7] .

India's TB-control response must balance two essential objectives: protecting the population from infection and protecting patients from discrimination, loss of confidentiality and denial of essential services[8] . A purely coercive approach may increase stigma and discourage individuals from seeking care, whereas a rights-based public-health framework can promote trust, treatment adherence and community cooperation[9] . Legal measures should consequently be evidence-based, proportionate, non-discriminatory and supported by accessible healthcare services.—Although India set an ambitious goal of TB elimination before the World Health Organization's global 2030 target, persistent challenges—including inadequate diagnostic capacity, shortages of health personnel, treatment interruptions, multidrug-resistant TB, malnutrition, stigma and uneven implementation—continue to hinder progress[10] . In this context, examining public health laws for the prevention and control of tuberculosis in India is essential[11] . Such an examination can clarify the legal basis of TB surveillance and notification, the responsibilities of public authorities and healthcare institutions, the relationship between public-health protection and individual rights, and the need for coordinated, community-centred

implementation. Strengthening the legal framework alongside NTEP interventions may contribute to earlier detection, improved treatment outcomes, reduced transmission and more equitable progress towards TB elimination[12] .


1. Tuberculosis as a Public Health Problem

Mycobacterium tuberculosis causes Tuberculosis (TB), which acts as a primary infectious illness and stands among the biggest global public health hurdles. Because transmission happens mainly via air when infected individuals cough, speak, sneeze, or release airborne droplets, managing it demands synchronized public health actions[13] . In contrast to ailments hitting just single people, TB can move between households, neighborhoods, clinics, job sites, and dense areas alike. Thus, successful TB management needs suitable clinical care plus organized steps for spotting cases early, reporting them, tracing contacts, stopping spread, and treating those carrying the bug or having active sickness[14] .



2. The Public Health Strategy for Stopping and Managing Tuberculosis.

Stopping and managing tuberculosis requires mixing together finding cases, diagnosing them, treating them, tracking contacts, watching trends, preventing spread, controlling infections, and teaching people about health. Finding those who carry contagious TB and making sure they get right care on time is key to breaking how it spreads. Also, people who met sick TB patients must be found, checked out, and given stop-treatment if needed[15] . Checking and handling people at higher risk for catching TB can lower the chance of moving from sleeping infection to real sickness. These actions show that fighting TB needs more than just treating patients clinically. It demands a public health strategy covering the whole population, bringing together state health officials, hospitals, labs, doctors, local groups, and various other partners involved[16] .



3. Tuberculosis Control in the Indian Context

India bears a significant load from tuberculosis, meaning that stopping and managing it forms a key part of its national public health setup. How big this problem gets depends on many things such as money issues, poor diet, too many people in small spaces, late finding of sickness, hard time getting care, shame and unfair treatment, plus problems with medicine courses. New kinds of bugs that fight drugs and how to handle them make controlling tuberculosis even harder now[17]. Currently, tuberculosis management within India has gradually moved away from focusing mainly on treating the illness itself toward adopting a broader strategy that highlights early detection, ensuring everyone can get care, reporting cases, stopping spread, helping patients, watching trends, and getting communities involved. Having both government-run and private medical groups take part matters greatly since people suffering from TB might go to various sections of the health network for their tests and medicines[18].

4. Necessity of Sanitary Regulations Regarding TB Management.

Because tuberculosis spreads between people, there is a requirement for rules and regulations that back up efforts made by public health officials. Laws concerning public health give official power and a structure so that state leaders may carry out watching for sickness, sending reports, controlling how medical care works, guarding the well-being of everyone, and putting into action plans designed to stop the spread of illness[19]. However, applying legal tools within tuberculosis management requires keeping a proper equilibrium between safeguarding community well-being and securing personal liberties. Actions designed to stop spread need execution that honors private life, secret-keeping, self-respect, fairness, and lack of bias. Such care matters greatly since shame tied to tuberculosis might stop people from getting checked or finishing their medicine courses[20].

5. Development of Sanitary Protocols and Statutory Actions Against Pulmonary Consumption.

Historically, efforts to manage tuberculosis developed parallel to growing scientific knowledge regarding the illness itself. Prior to the discovery of *Mycobacterium tuberculosis*, officials responsible for public health possessed restricted comprehension concerning how the infection spread, which resulted in having significantly

fewer practical tools available for managing the condition[21]. After the specific pathogen causing the disease was identified and mechanisms of transmission were better grasped, actions including isolating patients, monitoring cases, and providing care within institutions emerged as vital parts of broader strategies to control TB. Following the creation of successful anti-tuberculosis drug regimens, managing the illness changed greatly since long-term hospital stays became less important and therapy could happen mostly outside clinics[22]. Thus, current tuberculosis management focuses on spotting cases quickly, curing them well, stopping spread, watching trends, finding contacts, and caring for patients individually. This past shift shows clearly that laws and rules for public health need to adjust when science and medical methods evolve[23].

6. The Legal Structure and Management of Public Wellness Within India.

In India, efforts regarding TB prevention and control function inside a wider structure made up of constitutional rules, laws on public health, official policies, specific regulations, court decisions, and country-wide programs[24]. This legal setup matters greatly for setting duties connected to reporting diseases, watching over cases, providing medical care, and shielding people hit by tuberculosis. Simultaneously, managing tuberculosis requires engagement from various actors such as national and regional administrations, public health bodies, medical practitioners, diagnostic facilities, independent care centers, and non-governmental groups[25]. Successful execution thus relies not just upon having statutes but equally on synchronization, organizational strength, rule application, knowledge levels, and how reachable medical support remains[26].

7. Justification for Examining Public Health Statutes Regarding Tuberculosis Management Within India.

Notwithstanding major progress regarding detection, therapy, and community-level actions, tuberculosis remains a difficult problem within India. Problems including slow identification, broken courses of medicine, resistant strains, social shame, privacy concerns, unfair treatment, and alignment efforts among state and non-state providers highlight why one must look closely at the laws that manage stopping and managing the disease[27]. Consequently, examining public health statutes concerning tuberculosis holds importance since it aids evaluation if current rules sufficiently back disease control while protecting rights and dignity of those impacted. This review may likewise reveal holes in execution and offer grounds for bolstering India's statutory and public health reaction against tuberculosis[28].

8. Research Gap and the Current Study's Justification

The legal aspects of TB prevention and control need careful consideration, despite the fact that the disease has been thoroughly researched from a medical and epidemiological standpoint. Important legal and ethical issues are raised, in particular, by the connection between public health goals, required disease notification, surveillance, patient confidentiality, nondiscrimination, treatment compliance, and individual rights[29].

Therefore, the current study aims to investigate the public health laws and associated legal mechanisms governing the prevention and control of tuberculosis in India, paying special attention to their efficacy, implementation difficulties, and compatibility with individual rights and public health goals[30] .

Aim of the Study

In order to understand and analyse the legal and regulatory regime relating to the prevention and control of tuberculosis in India with a special focus on their efficacy, implementation and individual rights protection.

Objectives of the Study

To study the legal and regulatory regime relating to the prevention and control of tuberculosis in India.

To analyse the importance of the mandate for TB Notification and Surveillance for the control of disease.

To analyse the obligations of public and private sector hospitals and medical practitioners with regard to TB control within the current regime.

To analyse the issue of public health protection and individual rights within the regime of TB control.

To identify the key challenges that exist within the legal and implementation regime for TB control.

To analyse the role of NTEP and other regulatory measures for meeting the objectives of TB-control.

To recommend measures for improving the legal regime for TB control in India.

3. THE HISTORICAL DEVELOPMENT OF TUBERCULOSIS PREVENTION AND CONTROL IN INDIA

3.1 Introduction

Tuberculosis has been one of the most significant public health problems of India in recent decades. The strategy related to tuberculosis prevention and control has been developing over the years due to changes in science, diagnostic procedures, medical infrastructure, and public health approaches. The Indian experience of TB prevention and control can be characterized by the development from rather limited and treatment-focused measures to comprehensive measures, which include early diagnostics, standard treatment, monitoring, prevention, community involvement and cooperation between different sectors.

The historical development of TB prevention and control in India is essential for analyzing the contemporary public health system. The transition from the National Tuberculosis Program to the Revised National Tuberculosis Control Program and further to the National Tuberculosis Elimination Program reveals changes in priorities and strategies of the country in dealing with tuberculosis. The existing literature underlines the significance of preventive measures and more extensive approaches such as One Health approach.

3.2 Initial Measures for Tuberculosis Control in India

The initial efforts made for the control of tuberculosis in India primarily emphasized on patient detection and

treatment, and disease prevention through vaccines and other public health measures. The early measures for tuberculosis control in India were hampered by poor healthcare infrastructure, shortage of diagnosis facilities, and the absence of any systematic surveillance.

BCG vaccine programme was one of the many early preventative interventions used in India. But the increasing prevalence of tuberculosis proved that the use of vaccination alone was not enough to control the disease. Hence there was an urgent need for a more planned and programmatic approach to prevent and treat tuberculosis.

3.3 Formation of National Tuberculosis Programme (NTP)

One of the significant events in the history of tuberculosis control in India was the formation of National Tuberculosis Program (NTP) in 1962. The program attempted to deliver organized tuberculosis services through the available public healthcare network.

The National Tuberculosis Program primarily consisted of the following:

Detection of tuberculosis cases.

Diagnosis using the available laboratories.

Anti-tuberculosis treatment.

Expansion of TB services through public health care system.

Integration of TB activities with healthcare services.

However, there were certain problems associated with this programme. Inadequate case finding, intermittent treatment, poor compliance and inadequate monitoring were some of the problems with this programme. Consequently, the impact of this programme remained limited, and tuberculosis continued to remain a public health challenge.

These deficiencies of the NTP programme showed the need for a well-structured and monitored standard TB control programme.

3.4 Implementation of the Revised National Tuberculosis Control Programme (RNTCP)

In order to address the deficiencies associated with the earlier programme, the Revised National Tuberculosis Control Programme (RNTCP) was launched in India. The RNTCP was formulated using the internationally known DOTS (Directly Observed Treatment, Short-course) strategy.

Implementation of the RNTCP was one of the major breakthroughs in India in terms of TB control. It aimed at ensuring proper diagnosis and completion of the treatment.

The key elements of the DOTS strategy were as follows:

Political and administrative commitment.

Quality assured diagnosis.

Short course chemotherapy.

Direct observation of treatment if necessary.

Recording and reporting.

However, through the implementation of the RNTCP, the process of monitoring and reporting of the disease as well as the way patients adhered to treatment were improved.

3.5 Expanding TB Control Strategies

In the course of time, the Indian strategy of TB control expanded beyond just diagnosing and treating TB. The program acknowledged the need for multi-faceted interventions in order to ensure TB control.

These included:

Early and proper diagnosis of the disease.

Access to treatment.

Adherence to treatment and support to the patient.

Management of drug resistant TB.

Monitoring.

Private health sector involvement.

Community participation.

Prevention of transmission of TB.

Thus, the strategy went from being a disease specific treatment to a public health strategy. The increased complexity of the disease, in particular, the rise of multidrug resistant TB proved the need for advanced diagnostic facilities and improved treatment and monitoring strategies.

3.6 Shift from RNTCP to the National Tuberculosis Elimination Programme

New epidemiological trends in TB led to a further shift in India's TB control strategy. Thus, the RNTCP was rebranded as the National Tuberculosis Elimination Programme (NTEP).

The name change signified a new set of objectives for the programme. Rather than aiming at the control of TB, the programme now aimed at its elimination as a public health issue.

Key features of the NTEP include:

Early detection and access to TB care for all.

Early start of treatment.

Prevention and treatment of drug-resistant TB.

Tuberculosis preventive therapy.

Contact tracing.

Case finding.

Surveillance and reporting.

Private sector engagement.

Patient-centred care.

Community involvement.

Hence, the move to NTEP signifies a shift from the treatment-focused approach to TB control to an elimination approach.

3.7 Enhancing Preventive Measures

One important development in the history of TB control is the growing focus on prevention strategies. In the old days, TB control was mainly about identifying and treating active TB cases. However, the new methods have understood that prevention and not just treatment is necessary to eliminate TB.

Preventive measures include:

Identification of infectious TB cases early on.

Preventing the spread of infection within families and communities.

Contact tracing.

Identification of people who are at a higher risk of TB.

Giving TB preventive therapy where possible.

Infection prevention and control.

Education.

Better nutrition and social services.

Another fact emerging from the literature on tuberculosis prevention is the influence of environmental factors on TB. Thus, effective prevention must go beyond the field of healthcare.

3.8 One Health Approach

One Health approach gives a wider framework for TB prevention and control. It focuses on the relationship between human health, animal health and the environment.

The One Health approach is especially useful when dealing with zoonotic TB and other scenarios where the interaction between people, animals and the environment can play a role in the spread of disease.

This approach involves:

Public health officials.

Medical experts.

Veterinarians.

Environmental agencies.

Researchers

Communities

3.9 Transition from Control of Disease to Patient-Centeredness

The third noteworthy shift in India's strategy for TB control has been the growing appreciation of the requirement for a patient-centered approach.

The problems faced by patients suffering from tuberculosis are:

Social stigma.

Economic constraints.

Unemployment or loss of income.

Accessibility to medical facilities.

Treatment is prolonged.

Side effects of drugs.

Psychological and social issues.

Thus, for effective TB control, it is essential not only to provide the medicine but also to ensure that patients have access to treatment.

Patient-centeredness emphasizes treatment compliance, social and nutritional support, removing barriers to healthcare and engaging the community.

3.10 The Role of Surveillance and Monitoring

Surveillance and monitoring have gained importance as an integral part of the modern TB control programme. The early programmes encountered problems due to the lack of knowledge about the disease occurrence and its treatment results.

An efficient surveillance system will help health authorities to:

- Determine TB patients.

- Evaluate disease patterns.

- Evaluate treatment results.

- Find out the deficiencies in health care services.

- Find out high disease burden areas.

- Plan intervention measures.

Thus, the evolution of the TB control programme also included gradual improvements in the recording, reporting, notification, and monitoring systems.

Such methods are especially important in terms of health governance as they allow health authorities to organize disease control activities and evaluate the impact of interventions.

3.11 Main Obstacles in India's TB Control

In spite of the considerable progress, India is still encountering some problems in relation to TB control and elimination.

Among the main problems there are:

- A heavy burden of the disease.

- Late diagnosis.

- Undetected/unreported cases.

- Drug-resistant TB.

- Interruption/non-compliance with the treatment regimen.

- Poverty and malnutrition.

- Overcrowdedness.

- Stigma and discrimination.

- Limited awareness.

- Accessibility problems.

- Coordination of public and private health care.

6 CONCLUSION

Tuberculosis remains a major public health issue in India, but considerable advances have been achieved in preventing and controlling it. The evolution from the BCG vaccination and the National Tuberculosis Programme (NTP) to the Revised National TB Control Programme (RNTCP) and the National TB Elimination Programme

(NTEP) illustrates the ongoing development of India's tuberculosis control system. The three studies show that in order to eliminate TB a thorough and coordinated approach is needed. For the purpose of reducing transmission it is essential to carry out early diagnosis, provide effective and complete treatment, trace contacts, offer preventive treatment, maintain surveillance and implement infection control. Nevertheless, problems such as poverty, malnutrition, overcrowding, stigma, migration, inadequate access to healthcare and drug resistance still have an impact on TB control efforts. The One Health approach also extends the prevention of tuberculosis by acknowledging the link between human health, animal health and environmental conditions, so cooperation between various sectors is needed if both human and zoonotic forms of tuberculosis are to be effectively dealt with. The literature surveyed suggests that it will not be possible to eliminate tuberculosis in India by relying solely on medical treatment. It is necessary for strong public health policies, effective notification and surveillance, collaboration between the public and private sectors, the involvement of the community, and consideration of social determinants all to play their part. Moreover, TB control measures should ensure that patients' privacy, dignity and rights are protected. It follows that an integrated, people-centred and multisectoral approach can thus form a more solid basis for attaining sustainable TB control and eventual elimination in India.

REFERENCES

- [1] D. Zenner et al., "State of the global tuberculosis epidemic: burden, progress towards WHO End TB Strategy targets and systemic challenges," *Lancet Microbe*, p. 101433, Jul. 2026, doi: 10.1016/j.lanmic.2026.101433.
- [2] A. Mundra, P. Kothekar, P. Deshmukh, and A. Dongre, "Why tuberculosis patients under revised national tuberculosis control programme delay in health-care seeking? A mixed-methods research from Wardha District, Maharashtra," *Indian J. Public Health*, vol. 63, no. 2, p. 94, 2019, doi: 10.4103/ijph.IJPH_138_18.
- [3] V. K. Arora and K. K. Chopra, "Journey of tuberculosis control in India," *Indian Journal of Tuberculosis*, vol. 66, no. 1, pp. 178–183, Jan. 2019, doi: 10.1016/j.ijtb.2019.02.002.
- [4] L. S. Chauhan and J. Tonsing, "Revised national TB control programme in India," *Tuberculosis*, vol. 85, no. 5–6, pp. 271–276, Sep. 2005, doi: 10.1016/j.tube.2005.08.003.
- [5] M. Najafizada, A. Rahman, Q. Taufique, and A. Sarkar, "Social determinants of multidrug-resistant tuberculosis: A scoping review and research gaps," *Indian Journal of Tuberculosis*, vol. 68, no. 1, pp. 99–105, Jan. 2021, doi: 10.1016/j.ijtb.2020.09.016.
- [6] M. M. Kavanagh, L. O. Gostin, and J. Stephens, "Tuberculosis, human rights, and law reform: Addressing the lack of progress in the global tuberculosis response," *PLoS Med.*, vol. 17, no. 10, p. e1003324, Oct. 2020, doi: 10.1371/journal.pmed.1003324.

- [7] L. Chauhan, "Public health in India: Issues and Challenges," *Indian J. Public Health*, vol. 55, no. 2, p. 88, 2011, doi: 10.4103/0019-557X.85237.
- [8] N. Giri et al., "Balancing Public Health Priorities and individual rights in community tuberculosis control," *Indian Journal of Tuberculosis*, vol. 73, pp. S30–S35, Jun. 2026, doi: 10.1016/j.ijtb.2026.02.021.
- [9] A. Chakrabartty, P. Basu, K. M. Ali, and D. Ghosh, "Tuberculosis related stigma attached to the adherence of Directly Observed Treatment Short Course (DOTS) in West Bengal, India," *Indian Journal of Tuberculosis*, vol. 66, no. 2, pp. 259–265, Apr. 2019, doi: 10.1016/j.ijtb.2019.03.005.
- [10] A. Poddar, S. Mukherjee, and S. R. Rao, "Enhancing India's Health Security Efforts Against Mycobacterium Tuberculosis : Gaps and Opportunities," *Health Secur.*, vol. 23, no. 4, pp. 230–241, Aug. 2025, doi: 10.1089/hs.2024.0109.
- [11] S. Chowdhury, G. Phutke, S. Patil, and Y. Jain, "The ethics of compulsory notification of tuberculosis," *Indian J. Med. Ethics*, pp. 01–03, Mar. 2019, doi: 10.20529/IJME.2019.013.
- [12] N. Gawde, "Do we need notification of tuberculosis? A public health perspective," *Indian J. Med. Ethics*, no. 1, Jan. 2013, doi: 10.20529/IJME.2013.014.
- [13] M. Sarkar and J. Sarkar, "Transmission of Mycobacterium Tuberculosis," *Journal of The Association of Physicians of India*, vol. 73, no. 9, pp. 91–96, Sep. 2025, doi: 10.59556/japi.73.1113.
- [14] L. Martinez, Y. Shen, E. Mupere, A. Kizza, P. C. Hill, and C. C. Whalen, "Transmission of Mycobacterium Tuberculosis in Households and the Community: A Systematic Review and Meta-Analysis," *Am. J. Epidemiol.*, vol. 185, no. 12, pp. 1327–1339, Jun. 2017, doi: 10.1093/aje/kwx025.
- [15] T. Garg, L. H. Chaisson, F. Naufal, A. E. Shapiro, and J. E. Golub, "A systematic review and meta-analysis of active case finding for tuberculosis in India," *The Lancet Regional Health - Southeast Asia*, vol. 7, p. 100076, Dec. 2022, doi: 10.1016/j.lansea.2022.100076.
- [16] N. Sharma, S. Basu, and K. K. Chopra, "Achieving TB elimination in India: The role of latent TB management," *Indian Journal of Tuberculosis*, vol. 66, no. 1, pp. 30–33, Jan. 2019, doi: 10.1016/j.ijtb.2018.10.006.
- [17] A. Bhargava, M. Bhargava, and A. Juneja, "Social determinants of tuberculosis: context, framework, and the way forward to ending TB in India," *Expert Rev. Respir. Med.*, vol. 15, no. 7, pp. 867–883, Jul. 2021, doi: 10.1080/17476348.2021.1832469.
- [18] K. S. Sachdeva et al., "Paradigm shift in efforts to end TB by 2025," *Indian Journal of Tuberculosis*, vol. 67, no. 4, pp. S48–S60, Dec. 2020, doi: 10.1016/j.ijtb.2020.11.001.
- [19] S. Chowdhury, G. Phutke, S. Patil, and Y. Jain, "The ethics of compulsory notification of tuberculosis," *Indian J. Med. Ethics*, pp. 01–03, Mar. 2019, doi: 10.20529/IJME.2019.013.
- [20] K. Shringarpure et al., "Patient adherence to tuberculosis treatment in the Indian subcontinent: systematic review and meta-synthesis of qualitative research," *BMJ Open*, vol. 13, no. 5, p. e063926, May 2023, doi: 10.1136/bmjopen-2022-063926.
- [21] P. C. Karakousis and G. Mooney, "Respiratory Isolation for Tuberculosis: A Historical Perspective," *J. Infect. Dis.*, vol. 231, no. 1, pp. 3–9, Feb. 2025, doi: 10.1093/infdis/jiae477.
- [22] E. D. Zwick and C. S. Pepperell, "Tuberculosis sanatorium treatment at the advent of the chemotherapy era," *BMC Infect. Dis.*, vol. 20, no. 1, p. 831, Dec. 2020, doi: 10.1186/s12879-020-05539-w.
- [23] J. Gittler, "Controlling Resurgent Tuberculosis: Public Health Agencies, Public Policy, and Law," *J. Health Polit. Policy Law*, vol. 19, no. 1, pp. 107–147, Feb. 1994, doi: 10.1215/03616878-19-1-107.
- [24] A. Khanna, R. Saha, and N. Ahmad, "National TB elimination programme - What has changed," *Indian J. Med. Microbiol.*, vol. 42, pp. 103–107, Mar. 2023, doi: 10.1016/j.ijmm.2022.10.008.
- [25] R. P. Suseela and M. Shannawaz, "Engaging the Private Health Service Delivery Sector for TB Care in India—Miles to Go!," *Trop. Med. Infect. Dis.*, vol. 8, no. 5, p. 265, May 2023, doi: 10.3390/tropicalmed8050265.
- [26] R. PS, M. Shannawaz, M. E. Mathew, and K. S. Sachdeva, "Facilitators and Barriers for Private Health Sector Engagement for TB Care in India: A Systematic Review and Meta-Synthesis of Qualitative Research," *Glob. Health Sci. Pract.*, vol. 12, no. 4, p. e2400034, Aug. 2024, doi: 10.9745/GHSP-D-24-00034.
- [27] C. T. Sreeramareddy, Z. Z. Qin, S. Satyanarayana, R. Subbaraman, and M. Pai, "Delays in diagnosis and treatment of pulmonary tuberculosis in India: a systematic review," *The International Journal of Tuberculosis and Lung Disease*, vol. 18, no. 3, pp. 255–266, Mar. 2014, doi: 10.5588/ijtld.13.0585.
- [28] S. A. Nair et al., "Standards for TB care in India: A tool for universal access to TB care," *Indian Journal of Tuberculosis*, vol. 62, no. 4, pp. 200–206, Oct. 2015, doi: 10.1016/j.ijtb.2015.11.004.
- [29] A. Patange et al., "Informed consent and confidentiality in community-level tuberculosis interventions," *Indian Journal of Tuberculosis*, vol. 73, pp. S52–S57, Jun. 2026, doi: 10.1016/j.ijtb.2026.02.015.
- [30] Y. Lee, "Future directions for notifiable diseases: tuberculosis-related laws in the Philippines," *Global. Health*, vol. 14, no. 1, p. 87, Dec. 2018, doi: 10.1186/s12992-018-0405-2.